

The effect of geomagnetic storms on suicide

Charmaine Gordon¹, Michael Berk²

1. Department of Psychiatry, Faculty of Health Science, University of the Witwatersrand Medical School,

2. Department of Psychiatry, University of Melbourne, Geelong, Victoria, Australia

ABSTRACT

Objectives: To correlate geomagnetic storm activity with suicide rates. **Design:** A retrospective analysis over a 13 year period, January 1980 to December 1992. **Setting:** Hermanus Magnetic Observatory (data on geomagnetic storm activity), South African Central Statistical Services (data on suicide rates). **Subjects:** Nil. **Outcome measures:** Geomagnetic storm activity and suicide rates. **Results:** A significant correlation ($r=0,6964$; $p<0,01$) was found between the mean total of suicides and the mean average of storm activity during the same period. This correlation was true of both male ($r=0,6301$; $p<0,025$) and female ($r=0,7544$; $p<0,005$) suicides. **Conclusions:** Geomagnetic storm activity is correlated with suicide, and confirms previous research suggesting an impact of ambient magnetic field activity on behaviour.

Keywords: Suicide, Geomagnetic, Depression, Behaviour, Magnetic

Introduction

The influence of magnetism on the human body and behaviour is not a new concept. Franz Mesmer (1734-18-15), a student of theology, astrology and medicine, had a basic theory that an invisible magnetic fluid emanating from the stars influenced the health of human beings. An imbalance of this fluid in the body causes illness and could be rectified by the touch of a magnetic wand. Later Mesmer became convinced that his hands had possessed magnetic power and that the magnetic fluid also flowed from his own body in the form of 'animal magnetism.'

Since the turn of this century empirical studies regarding geomagnetic influences on humans were described.^{1,2,3} Kay (1994), found a high correlation between the admission rate of 3449 patients diagnosed as suffering from depression, to a psychiatric hospital in Britain, and geomagnetic data collected over a 10 year period.⁴ The majority of studies have centred around the effect of temperature changes on behaviour⁵, and on the influence of man-made industrial and domestic electro-magnetic fields on humans and other living systems.^{6,7} Industrial and domestic electro-magnetic fields are different from biogeomagnetism, which is the study of possible low-frequency electromagnetic field disturbances from the solar terrestrial environment on humans and other living systems.⁸ The main cause of geomagnetic storms is a sudden in-

crease in charged particles emanating from solar flares. These storms can last several hours and are a global phenomenon. Geomagnetic storms follow a seasonal pattern, are most numerous during the equinoxes.⁹

As far as natural geomagnetism is concerned, it is documented that electric storms cause homing pigeons to lose their sense of direction.¹⁰ It is also known that bees¹¹, bacteria and snails orientate themselves to the natural geomagnetic flow of the earth, different in the opposite hemispheres.¹² In all these species, magnetite, a magnetic crystal, can be found.¹³

Numerous laboratory studies have shown the effects of magnetic fields on living organisms.^{14,15} Neurobiochemical studies have shown that energy fields at the same strength as geomagnetic field strengths (GMF), decreases the hydrophobic character of the cell membrane leading to changing permeability.^{11,16,17} It has been shown that magnetic fields alter the electrical properties of solutions and their physiological effects.¹⁸ Membrane potential and electrical activity of neurons was shown to be altered when normal physiological solution was replaced by CaCl_2 which was exposed to a magnetic field. Frequency dependence of electromagnetic field-induced Calcium-ion efflux from chicken brain tissues have been shown.¹⁹

There are indications that the pineal gland, which regulates the circadian rhythm and which is involved with the regulation and production of melatonin²⁰, is a magnetosensitive system.²¹ The nightly production and secretion of melatonin by the pineal gland provides information regarding the time-of-day and time-of-year to the rest of the body, thus acting as a time-cued or 'Zeitgeber' function.²² In mammals the circadian rhythm of melatonin is synchronised by the prevailing light: dark environment with

Correspondence:

Prof M Berk, Dept of Psychiatry, University of Melbourne, Swanston Centre, PO Box 281, Geelong, Victoria, 3220, Australia.
email: mikebe@barwonhealth.org.au

the retina of the eyes doing the photoreception required for the induction of this rhythm.²² Besides visible light, certain ultra violet wavelengths as well as extremely low frequency electric and magnetic fields may be involved with the melatonin rhythm.²² Circadian abnormalities may result from a failure in the entrainment processes of internal clocks by environmental time cues.^{23,24} It has been demonstrated that individuals placed in an underground bunker, thus shielded from the ambient electromagnetic field, showed gradual desynchronization and lengthening of circadian rhythms, which could be re-synchronized when exposed to artificial magnetic fields^{25,26,27}

According to Kay⁴, storms in spring would enhance the suppressing effect of increasing daylight on melatonin synthesis, leading to a phase advance in the circadian rhythm, while the effect of storms in autumn would tend to be partially compensated for by the pineal response to decreasing light intensity. The pineal gland was shown to be sensitive to changes in magnetic field exposure in laboratory animals.²¹ Alteration of the direction of the geomagnetic field is associated with a reduction of spontaneous electrical activity in the pineal gland, induction of ultrastructural changes in the pineal gland, reduction of N-acetyltransferase and hydroxyindole-O-methyltransferase, reduction of nocturnal cyclic adenosine 3,5-monophosphate content in the pineal gland, inhibition of melatonin secretion, increase glucose uptake in the pineal gland, inhibition of rhythmicity of responses of Purkinje cells to melatonin in the cerebellum and to inhibition of the enhanced nocturnal analgesic response to morphine in mice.²¹

Considering the earth's mainly metallic molten core¹³ and since the daily heating and cooling of the atmosphere displaces the jet streams north and south, a more or less regular circadian variation in the magnetic field of the earth can be observed on the ground.²⁸ This could imply that the natural circadian rhythmicity of the earth's magnetic field could have a 'Zeitgeber' or time cued function.²⁹ However, it is possible that this natural magnetic flow of the earth, can be disturbed by geomagnetic storms.¹³

Geomagnetic storms are mainly caused by a sudden increase in charged high-energy particles and clouds of ionised gas or plasma, emanating from solar flares.³⁰ Storms are a global phenomena and seem to follow a seasonal pattern, being most numerous during the equinoxes.³¹ Storms have been found to desynchronise ambulatory behaviour patterns in rats³², and in mice are associated with decreased nocturnal analgesic effects of morphine, possibly mediated via the pineal gland.³³

While a correlation between Seasonal Affective Disorder and suicide in South Africa has been described (Kerr, unpublished data) the effect of geomagnetic influence on suicide still need to be determined. Furthermore, studies correlating geomagnetic influence on depression and suicide still lack in the Southern hemisphere. This study investigates whether the equinoxal increase in geomagnetic storms will correlate with the bimodal annual distribution of incidence of suicide in a South African population..

Method

Dates of all geomagnetic storms during a 13 year period between January 1980 and December 1992 was obtained from the Hermanus Magnetic Observatory, which is representative for the region from which the suicide data was collected, namely all of South Africa.

K indices from this observatory on the days of the geo-


magnetic storms were used. The K indices are quasi-logarithmic 3-hour means of geomagnetic disturbances on any particular day. Three magnetic field components are continually being registered by the observatory. D, which represents the magnetic declination, H, the horizontal component and Z the vertical component. As it is only necessary to determine size of the disturbance in this study, the Z component was used for correlation, seeing that this component contributes more than 90% to the calculation of the total magnetic field strength.

National registered suicide mortality data for the same period, for the whole South African population, was obtained from the South African Central Statistic Services (SACSS). An autopsy performed by a district surgeon is required for all deaths with an unnatural cause. The autopsy report is sent to a magistrate. An inquest is then held, the main aim of which is to establish the cause of death. This cause is then forwarded to the SACSS. Causes of death are classified according to a system based on the ICD-9 (International Classification of Disease)³⁴, reference E950-959 and E979. Mortality monthly means were calculated in the same way as for the geomagnetic data.

Monthly means for every month was determined, and then a cumulative mean for that specific month over the 13 year period was calculated. This method was used as to eliminate a Type II error. The Pearson's Product Moment Correlation Coefficient method from the Statistix PC package was used to determine correlation.

Results

The relationship between suicide and geomagnetic storm activity is shown in Figure 1. The December and January periods do not show a correlation between geomagnetic storm activity and suicide. In a year by year breakdown of the data, this pattern was observed for every year during those months, suggesting that some other factors may be involved. Temperature may have an effect on depression and suicide.⁵ To control for this variant, data for these months have been omitted from the calculations. By calculating in this manner, a significant correlation ($r = 0,6964$; $p < 0,01$) was found between the mean total of suicides for the remaining ten months, and the mean average of storm activity during the same period. The correlation of male suicide rates separately for this ten month period yielded a significant correlation ($r = 0,6301$; $p < 0,025$). The number of female victims (287) was four



times less than that for males (1200). However, their correlation with geomagnetic storm activity was highly significant ($r = 0,7544$; $p < 0,005$).

Discussion

Changes in the environmental magnetic field, which are directly associated with solar activity as well as circadian variants in the earth's magnetic field strengths are capable of disturbing the electrical activity of the brain and could lead to various neuropsychiatric disturbances, such as SAD, depression and suicide. Due to the influence of magnetic fields on the pineal gland, it is possible that these effects are mediated through alterations in pineal melatonin functions.²¹ Phase advance in pineal circadian rhythms of melatonin synthesis may be a possible mechanism of causation, or be present as a consequence of 5-hydroxytryptamine or adrenergic system dysfunction associated with geomagnetic disturbances.⁴ Effects on cell membrane permeability, calcium channel activity and refined magneto-receptors are suggested as possible underlying biochemical mechanisms.⁴

Transcranial Magnetic Stimulation (TMS) is an experimental therapeutic modality. Experimental studies have shown that magnetic stimulation can cause 5HT_{1B} receptor downregulation. Like ECT, TMS can activate immediate early genes. Left Prefrontal TMS has been shown to increase ¹⁸FDG uptake with PET scanning. Antidepressant effects of left prefrontal TMS, at high (20Hz) frequency but not right prefrontal TMS has been reported. Younger, right handed patients are more likely to respond. Gabapentin may enhance the therapeutic response to TMS. Right prefrontal TMS may possibly be antimanic.^{35,36,37,38,39,40} Controlled data on short term use of TMS in recurrent depression is now available.⁴¹

From recent research it seems that violent versus non-violent suicide correlates differently with biometeorological influences, with violent suicides showing a stronger correlation.^{5,42} Furthermore, gender specific responses to geomagnetic disturbances could be expected as the pineal gland is involved with regulating sexual activities⁴³, as well as the fact that it plays a role in the varying melatonin levels involved in the menstrual cycle.⁴⁴

It is well documented that temperature has an effect on depression and suicide.⁵ As the December / January period is a period characterised by higher temperatures in South Africa, it is possible that this could play a role. Some studies further reports more attempted suicides in the last six months of the year, especially in December.^{45,46,47} This study controlled for this variant in a manner reported in other studies.⁴ This study did not, however, directly control for the effects of season. Flisher et al (1997) described a seasonal pattern in suicide in South Africa, with a peak in the spring (September/October) or summer and a trough in winter.⁴⁸ This pattern was described as being more pronounced for a sub-group that is less urbanized and for another sub-group with a relatively low standard of living.

Conclusion

The results of this study support the hypothesis that a geomagnetic storm activity is associated with suicide rates. This finding was observed in both male and female populations in South Africa. A weakness of this study was that although very accurate and complete data for the geomagnetic part of the

study exists, only monthly averages for the corresponding suicide rates could be obtained. Research using daily or weekly suicide figures may yield a tighter correlation. Further investigation to determine possible correlation between geomagnetic storm influence on violent versus non-violent suicides in both male and female populations could also to be undertaken.

References

- 1 Durkheim E. *Le Suicide*. Paris, 1897.
- 2 Dull T, Dull B. *Zusammenhänge zwischen Störungen des Erdmagnetismus und Häufungen von Todesfällen*. *Deutsche Medizinische Wochenschrift* 1935; 61 : 95.
- 3 Friedman H, Becker RO, Bachman CH. (*Geomagnetic parameters and psychiatric hospital admissions*. *Nature* 1963; 200 : 626-628.
- 4 Kay RW. *Geomagnetic storms : Association with Incidence of Depression as Measured by Hospital Admission*. *British Journal of Psychiatry* 1994; 164 : 403-409.
- 5 Maes M, De Meyer F, Thompson P, Peeters D, Cosyns P. *Synchronized annual rhythms in violent suicide rates, ambient temperature and the light-dark span*. *Acta Psychiatrica Scandinavica* 1994; 90 : 391-396.
- 6 World Health Organization. *Extremely Low Frequency Fields (ELF)*. Environmental Health Organization. 35. Geneva, 1984.
- 7 Hendee WR, Boteler JC. *The question of health effects from exposure to electromagnetic fields*. *Health Physics* 1994; 66 : 127-136.
- 8 Roederer JG. *Effects of natural magnetic field disturbances on biota: Fact or fiction. A report to the NRC Committee on Solar-Terrestrial Research*, 1995.
- 10 Beason RC, Nichols JE. *Magnetic orientation and magnetically sensitive material in a transequatorial migratory bird*. *Nature* 1984; 309 : 151
- 11 Adey WR. *Tissue interactions with non-ionizing electromagnetic fields*. *Physiological Review* 1981; 61 : 435-514.
- 12 Kirschvink JL, Jones DS, Mac Fadden BJ. (eds). *Magnetic Biomineralization and Magnetoreception in Organisms : A new Biomagnetism*. New York. Plenum Press 1985.
- 13 Parkinson WD. *Introduction to geomagnetism*. Edinburgh. Scottish Academic Press 1983.
- 14 Liboff AR, McLeod BRO. *Kinetics of channelized membrane ions in magnetic fields*. *Bioelectromagnetics* 1988; 9 : 39-51.
- 15 Lednev W. *Possible mechanisms for the influence of weak magnetic fields on biological systems*. *Bioelectromagnetics* 1991; 12 : 71-76.
- 16 Tenforde TS, Kaune WT. *Interaction of extremely low frequency electric and magnetic fields with humans*. *Health Physics* 1987; 53 : 585-606.
- 17 Marron MT, Goodman EM, Sharpe PT, et al. *Low frequency electric and magnetic fields have different effects on the cell surface*. *FEBS Letters* 1988; 230 : 13-16.
- 18 Ayrapetyan SN, Grigorian KV, Avanesian AS, Stamboltsian KV. *Magnetic fields alter electrical properties and their physiological effects*. *Bioelectromagnetics* 1994; 15 : 133-142.
- 19 Blackman CF, Benane SG, Elliott DJ, House DE, Pollock MM. *Influence of Electromagnetic Fields on the Efflux of Calcium Ions from Brain Tissue In Vitro: A Three-Model Analysis Consistent with the Frequency Response up to 510 Hz*. *Bioelectromagnetics* 1988; 9 : 215-227.
- 20 Cooper JR, Bloom FE, Roth RH. *The Biochemical Basis of Neuropharmacology*. New York. Oxford University Press 1974.

- 21 Sandyk R, Anninos PA, Tsagas N. Magnetic fields and seasonality of affective illness : implications for therapy. *International Journal of Neuroscience* 1991; 58 : 261-267.
- 22 Reiter RJ. Alterations of the circadian melatonin rhythm by the electromagnetic spectrum: a study in environmental toxicology. *Regulatory Toxicological Pharmacology* 1992;15 : 226-244.
- 23 Kripke DF, Mullaney D J, Atkinson M, Huey LY, Hubbart B. (1979). Circadian rhythm phases in affective illness. *Chronobiologia*. 6 : 4-12.
- 24 Souetre E, Salvati E, Pringuey D, Krebs B, Darcourt G. The circadian rhythm of plasma thyrotropin in depression and recovery. *Chronobiology international* 1986; 3 :197-205.
- 25 Wever R. Ueber die Beeinflussung der circadianen Periodik des Menschen durch schwache electromagnetische Felder. *Zeitschrift der Vergleichende Physiologie* 1967; 56 : 111-128.
- 26 Wever R. Einfluss schwacher elektro-magnetischer Felder auf die circadiane Periodik des Menschen. *Naturwissenschaften* 1968a; 55 : 29-32.
- 27 Wever R. Gestezmaessigkeiten der circadianen Periodik des Menschen geprüft an der Wirkung eines schwachen electrischen Wechselfeldes. *Pfluegers Archives* 1968b; 302 : 97-122.
- 28 Wiltschko W. The earth's magnetic field and bird orientation. *Trends in Neuroscience* 1980; 3 : 140-144.
- 29 Persinger MA, Ludwig HW, Ossenkamp P Psychophysiological effects of extremely low frequency of electromagnetic fields. A review. *Perception and Motor Skills* 1973; 36: 1131-1159.
- 30 Muir H. Watch out, here comes the sun. *New Scientist* 1996; 22-26.
- 31 Allen JH, Kihn EA. Major magnetic storms: Ap (1932-1989) and AA (1968-1988). Boulder, Colorado: National Geophysical Data Center 1990.
- 32 Persinger MA. Day time wheel running activity in laboratory rats following the geomagnetic event of 5-6 July 1974. *International Journal of Biometeorology* 1974; 20 : 19-22.
- 33 Ossenkopp KP, Kavaliers M, Hirst M. Reduced nocturnal morphine analgesia in mice following geomagnetic disturbance. *Neuroscience Letters* 1983; 40 : 321-325.
- 34 World Health Organization. *Manual of the international statistical classification of disease, injuries, and cause of death*. Geneva, 1978.
- 35 Massot O, Grimaldi B, Bailly JM, Kochanek M, Deschamps F, Lambrozo J, Fillion G. Magnetic field (MF) affects 5-HT_{1B} receptor function in brain: Molecular and cellular studies, *Eur Neuropsychopharmacol* 1988; 8 (Suppl 2): S119.
- 36 Belmaker RH, Grisar N. Transcranial magnetic stimulation in mania, *Eur Neuropsychopharmacol* 1988; 8, (Suppl 2): S120.
- 37 Schläpfer T E, Rupp F, Fisch HU, Hess C W. Effect of rTMS and ECT on gene expression: Same or different antidepressant mechanisms. *Eur Neuropsychopharmacol* 1998; 8 (Suppl 2): S121.
- 38 George MS, Bohning DE, Nahas Z, Shastri A, Lorberbaum JP, Speer AM, Vincent DJ, Roberts D, Stallings LE, Teneback C, Blumenthal K. Functional neuroimaging studies of TMS: SPECT, PET and fMRJ. *Eur Neuropsychopharmacol* 1998; 8 (Suppl 2): S122.
- 39 George MS, Wasserman EM, Williams WA, Callahan A, Ketter TA, Basser P, Hallet M, Post RM. Daily repetitive transcranial magnetic stimulation (rTMS) improves mood in depression. *Neuroreport* 1995; 6 : 1853-1856.
- 40 Pascual-Leone A, Keenan J, Freund S, Stinchfield Z, Tormos JM, Parker A, Ives J, Erb J, Birnbaum R. Repetitive transcranial magnetic stimulation trials in depression. *Eur Neuropsychopharmacol* 1988; 8 (Suppl 2): S123.
- 41 Klein E, Kreinin I, Chistyakov A, Koren D, Mecz L, Marmur S, Ben Shachar D, Feinsod M. Therapeutic efficacy of right prefrontal slow repetitive transcranial magnetic stimulation in major depression: a double-blind controlled study. *Arch Gen Psychiatry* 1999; 56: 315-320.
- 42 Maes M, Cosyns P, Meltzer HY, De Meyer F, Peeters D. Seasonality in Violent Suicide but not in Nonviolent Suicide or Homicide. *American Journal of Psychiatry* 1993; 150 (9) : 1380- 1385.
- 43 Tamarkin L, Baird CJ, Almeida OFX. Melatonin: a coordinating signal for mammalian reproduction. *Science* 1985; 277 : 714-715.
- 44 Brzezinski A, Lynch HJ, Seibel MM, et al. The circadian rhythm of plasma melatonin during the normal menstrual cycle and in amenorrheic woman. *Journal of Clinical Endocrinology and Metabolism* 1988; 66 : 891-895.
- 45 Whitlock A, Shapira K. Attempted suicide. *British Journal of Psychiatry* 1967; 113 : 423-434.
- 46 Schmidt C, Van Arsdol M. Completed and Attempted Suicides. *American Sociological Review* 1975; 20 : 273-283.
- 47 Wenz FV. Effects of season and sociological variables on suicidal behaviour. *Public Health Report* 1977; 92 : 233-239.
- 48 Flisher AJ, Parry CD, Bradshaw D, Juritz JM. Seasonal variation of suicide in South Africa. *Psychiatry Res* 1997; 15;66:13-22